arXiv:1401.3904v2 [math.FA] 17 Jan 2014

NORM ESTIMATES FOR SOLUTIONS OF ELLIPTIC BVPS OF THE DIRAC OPERATOR

DEJENIE A. LAKEW

Abstract. We present norm estimates for solutions of first and second order elliptic BVPs of the Dirac operator $D = \sum_{j=1}^{n} e_j \partial_{x_j}$ considered over bounded and smooth domain Ω of $\mathbb{R}^n$. The solutions whose norms to be estimated are in some Sobolev spaces $W^{k,p}(\Omega)$ and the boundary conditions as traces of solutions and their derivatives are in some Slobodeckij spaces $W^{\lambda,p}(\partial\Omega)$ where λ is some non integer but fractional number, for $1 \leq p < \infty$ and $k \in \mathbb{Z}$.

1. Algebraic and Analytic Rudiments of Cl_n

Let $\{e_j : j = 1, 2, ..., n\}$ be an orthonormal basis for $\mathbb{R}^n$ that is equipped with an inner product so that

$$e_i e_j + e_j e_i = -2\delta_{ij} e_0 \tag{1.1}$$

where δ_{ij} is the Kronecker delta. The inner product satisfies an anti commutative relation

$$x^2 = -\|x\|^2 \tag{1.2}$$

Therefore $\mathbb{R}^n$ with these properties of base vectors generates a non commutative algebra called Clifford algebra denoted by Cl_n.

The basis of Cl_n will then be

$$\{e_A : A \subset \{1 < 2 < 3 < ... < n\}\}$$

which implies:

$$\dim(Cl_n) = 2^n$$

The object e_0 used above is the identity element of the Clifford algebra Cl_n.

Representation of elemnets of Cl_n: every $a \in Cl_n$ is represented by

$$a = \sum e_A a_A \tag{1.3}$$

where a_A is a real number.

Date: December 30, 2013.

2000 *Mathematics Subject Classification.* Primary 46E35,47B38, Secondary 35C15.

Key words and phrases. Norm estimate, Dirac operator, trace, Sobolev spaces, Slobodeckij spaces, Elliptic BVP.

This paper is in final form and no version of it will be submitted for publication elsewhere.

Thus every $x = (x_1, x_2, ..., x_n) \in \mathbb{R}^n$ can be identified with $\sum_{j=1}^{n} e_j x_j$ of Cl_n and therefore we have an embedding

$$\mathbb{R}^n \hookrightarrow Cl_n$$

We also define what is called a Clifford conjugate of

$$a = \sum e_A a_A$$

as

$$\overline{a} = \sum \overline{e}_A a_A$$

where

$$\overline{e_{j_1}...e_{j_r}} = (-1)^r e_{j_r}...e_{j_1}$$

For instance for $i, j = 1, 2, ..., n$,

$$\overline{e}_j = -e_j, \quad e_j^2 = -1$$

and for

$$i \neq j : \overline{e_i e}_j = (-1)^2 e_j e_i = e_j e_i$$

Definition 1. *We define the Clifford norm of*

$$a = \sum e_A a_A \in Cl_n$$

by

$$\|a\| = \left((a\overline{a})_0\right)^{\frac{1}{2}} = \left(\sum_A a_A^2\right)^{\frac{1}{2}} \tag{1.4}$$

where $(a)_0$ *is the real part of* $a\overline{a}$.

The norm $\|.\|$ satisfies the inequality:

$$\|ab\| \leq c(n) \|a\| \|b\| \tag{1.5}$$

with $c(n)$ a dimensional constant.

Also each non zero element $x \in \mathbb{R}^n$ has an inverse given by :

$$x^{-1} = \frac{\overline{x}}{\|x\|^2} \tag{1.6}$$

$\lhd$ In the article it is always the case that $1 < p < \infty$ unless otherwise specified and Ω is a bounded and smooth (at least with C^1 - boundary $\partial\Omega$) domain of $\mathbb{R}^n$

A Clifford valued (Cl_n- valued) function f defined on Ω as

$$f : \Omega \longrightarrow Cl_n$$

has a representation

$$f = \sum_A e_A f_A \tag{1.7}$$

where $f_A : \Omega \longrightarrow \mathbb{R}$ is a real valued component or section of f.

Definition 2. *For a function $f \in C^1(\Omega) \cap C(\overline{\Omega})$, we define the Dirac derivative of f by*

$$Df(x) = \sum_{j=1}^{n} e_j \partial_{x_j} f(x) \tag{1.8}$$

A function $f : \Omega \longrightarrow Cl_n$ is called left monogenic or left Clifford analytic over Ω if

$$Df(x) = 0, \ \forall x \in \Omega$$

and likewise it is called right monogenic over Ω if

$$f(x)D = \sum_{j=1}^{n} \partial_{x_j} f(x) e_j = 0, \ \forall x \in \Omega$$

An example of both left and right monogenic function defined over $\mathbb{R}^n \backslash \{0\}$ is given by

$$\psi(x) = \frac{\overline{x}}{\omega_n \|x\|^n}$$

where ω_n is the surface area of the unit sphere in $\mathbb{R}^n$.

The function ψ is also a fundamental solution to the Dirac operator D and we define integral transforms as convolutions of ψ with functions of some function spaces below.

Definition 3. *Let $f \in C^1(\Omega, Cl_n) \cap C(\overline{\Omega})$.*
We define two integral transforms as follow:

$$\zeta_\Omega f(x) = \int_\Omega \psi(y - x) f(y) \, d\Omega_y, \quad x \in \Omega \tag{1.9}$$

$$\xi_{\partial\Omega} f(x) = \int_{\partial\Omega} \psi(y - x) \upsilon(y) f(y) \, d\partial\Omega_y, \quad x \notin \partial\Omega \tag{1.10}$$

The integral transform defined in (1.9) a domain integral is called the Theodorescu transform or the Cauchy transform. It is a convolution $\psi * f$ over Ω. The integral transform defined in (1.10) is some times called the *Feuter* transform as a boundary integral which again is a convolution $\psi * \upsilon f$ over $\partial\Omega$. $\upsilon(y)$ is a unit normal vector pointing outward at $y \in \partial\Omega$.

2. Sobolev and Slobodeckij Spaces

Definition 4. *For $1 < p < \infty$, $k \in \mathbb{N} \cup \{0\}$ we define:*

I: The Sobolev space $W^{k,p}(\Omega)$ as

$$W^{k,p}(\Omega) := \{f \in L^p(\Omega) : D^\alpha f \in L^p(\Omega), \ \|\alpha\| \leq k\}$$

with norm

$$\|f\|_{W^{k,p}(\Omega)}=\left(\sum_{\|\alpha\|\leq k}\int_{\Omega}|D^{\alpha}f|^{p}dx\right)^{\frac{1}{p}} \tag{2.1}$$

II: The Slobodeckij spaces for $0<\lambda<1$ as

$$W^{\lambda,p}(\partial\Omega):=\{f\in L^{p}(\partial\Omega):\int_{\partial\Omega}\int_{\partial\Omega}\frac{|f(x)-f(y)|^{p}}{|x-y|^{n+\lambda p-1}}d\sigma_{x}d_{\sigma y}<\infty\}$$

and norm is defined by

$$\|f\|_{W^{\lambda,p}(\partial\Omega)}=\left(\int_{\partial\Omega}\int_{\partial\Omega}\frac{|f(x)-f(y)|^{p}}{|x-y|^{n+\lambda p-1}}d\sigma_{x}d_{\sigma y}\right)^{\frac{1}{p}} \tag{2.2}$$

III: The Slobodeckij spaces for $\lambda=[\lambda]+\{\lambda\}$ where $0<\{\lambda\}<1$:

$$W^{\lambda,p}(\partial\Omega):=\{f\in W^{[\lambda],p}(\partial\Omega):\sum_{\|\alpha\|\leq[\lambda]}\int_{\partial\Omega}|Df|^{p}d\sigma_{x}+\sum_{\|\alpha\|=[\lambda]}\int_{\partial\Omega}\int_{\partial\Omega}\frac{|D^{\alpha}f(x)-D^{\alpha}f(y)|^{p}}{|x-y|^{n+\{\lambda\}p-1}}d\sigma_{x}d_{\sigma y}<\infty\}$$

and hence norm is given by

$$\|f\|_{W^{\lambda,p}(\partial\Omega)}=\left(\sum_{\|\alpha\|\leq[\lambda]}\int_{\partial\Omega}|Df|^{p}d\sigma_{x}+\sum_{\|\alpha\|=[\lambda]}\int_{\partial\Omega}\int_{\partial\Omega}\frac{|D^{\alpha}f(x)-D^{\alpha}f(y)|^{p}}{|x-y|^{n+\{\lambda\}p-1}}d\sigma_{x}d_{\sigma y}\right)^{\frac{1}{p}} \tag{2.3}$$

In the definitions of the Slobodeckij spaces and associated norms, the irregularity exponent $n+\{\lambda\}p-1$ is due to the fact that the dimension of $\partial\Omega$ is $n-1$ and $d\sigma$ is a hypersurface measure on $\partial\Omega$.

Slobodeckij spaces as subspaces of Sobolev spaces but with fractional exponents are analogues of the Hölder spaces in classical spaces of continuous functions.

3. Some Properties and Relations Between D,ζ_{Ω},τ and $\xi_{\partial\Omega}$

Proposition 1. *$D:W^{k,p}(\Omega,Cl_{n})\longrightarrow W^{k-1,p}(\Omega,Cl_{n})$ is continuous with*

$$\|Df\|_{W^{k-1,p}(\Omega,Cl_{n})}\leq\gamma\|f\|_{W^{k,p}(\Omega,Cl_{n})}$$

for $\gamma=\gamma(n,p,\Omega)$ a positive constant.

Proof. Let $f\in W^{k,p}(\Omega,Cl_{n})$. We need to show that

$$\|Df\|_{W^{k-1,p}(\Omega,Cl_{n})}\leq c\|f\|_{W^{k,p}(\Omega,Cl_{n})}$$

for some positive constant c.

$$f\in W^{k,p}(\Omega,Cl_{n})\Longrightarrow\|f\|_{W^{k,p}(\Omega,Cl_{n})}=\left(\sum_{\|\alpha\|\leq k}\int_{\Omega}|D^{\alpha}f|^{p}dx\right)^{\frac{1}{p}}<\infty$$

But then

$$\begin{aligned}
\|Df\|_{W^{k-1,p}(\Omega,Cl_n)} &= \left(\sum_{\|\alpha\|\leq k-1}\int_\Omega |D^\alpha f|^p dx\right)^{\frac{1}{p}} \\
&\leq \left(\sum_{\|\alpha\|\leq k-1}\int_\Omega |D^\alpha f|^p dx + \sum_{\|\alpha\|=k-1}\int_\Omega |D^\alpha f|^p dx\right)^{\frac{1}{p}} \\
&= \left(\sum_{\|\alpha\|\leq k}\int_\Omega |D^\alpha f|^p dx\right)^{\frac{1}{p}} \\
&= \|f\|_{W^{k,p}(\Omega,Cl_n)}
\end{aligned}$$

Therefore for $c = 1$, the proposition is proved. □

Proposition 2. *$D : L^p(\Omega) \longrightarrow W^{-1,p}(\Omega)$ is continuous for $1 < p < \infty$.*

Proof. Let $f \in L^p(\Omega)$. Then

$$\|Df\|_{W^{-1,p}(\Omega)} = \sup\{\frac{|\langle Df, v\rangle|}{\|v\|_{W_0^{1,q}(\Omega)}} : v \neq 0, v \in W_0^{1,q}(\Omega)\}$$

for $p^{-1} + q^{-1} = 1$.

But

$$|\langle Df, v\rangle| = |\langle f, Dv\rangle| \leq \|f\|_{L^p(\Omega)}\|Dv\|_{L^q(\Omega)} \leq \|f\|_{L^p(\Omega)}\|v\|_{W_0^{1,q}(\Omega)}$$

Thus by the Cauchy-Schwartz inequality we have

$$\frac{|\langle Df, v\rangle|}{\|v\|_{W_0^{1,q}(\Omega)}} \leq \frac{\|f\|_{L^p(\Omega)}\|v\|_{W_0^{1,q}(\Omega)}}{\|v\|_{W_0^{1,q}(\Omega)}} = \|f\|_{L^p(\Omega)}$$

Therefore

$$\begin{aligned}
\|Df\|_{W^{-1,p}(\Omega)} &= \sup\{\frac{|\langle Df, v\rangle|}{\|v\|_{W_0^{1,q}(\Omega)}} : v \neq 0,\ v \in W_0^{1,q}(\Omega)\} \\
&\leq \sup\{\frac{\|f\|_{L^p(\Omega)}\|v\|_{W_0^{1,q}(\Omega)}}{\|v\|_{W_0^{1,q}(\Omega)}} : v \neq 0,\ v \in W_0^{1,q}(\Omega)\} \\
&= \|f\|_{L^p(\Omega)}
\end{aligned}$$

□

Proposition 3. *(Mapping properties) ([4], [6])*

Let $k \in \mathbb{N} \cup \{0\}$ and $1 < p < \infty$. Then there are positive constants $\beta = \beta(n, p, \Omega)$, $\theta = \theta(n, p, \Omega)$ and $\delta = \delta(n, p, \Omega)$ such that

$$\zeta_\Omega : W^{k,p}(\Omega, Cl_n) \longrightarrow W^{k+1,p}(\Omega, Cl_n) \tag{3.1}$$

with

$$\|\zeta_\Omega f\|_{W^{k+1,p}(\Omega,Cl_n)} \leq \beta \|f\|_{W^{k,p}(\Omega,Cl_n)}$$

$$\xi_{\partial\Omega} : W^{\lambda,p}(\partial\Omega, Cl_n) \longrightarrow W^{\lambda+\frac{1}{p},p}(\Omega, Cl_n) \tag{3.2}$$

with

$$\|\xi_{\partial\Omega} f\|_{W^{\lambda+\frac{1}{p},p}(\Omega,Cl_n)} \leq \theta \|f\|_{W^{\lambda,p}(\partial\Omega,Cl_n)}$$

and

$$\tau : W^{k,p}(\Omega, Cl_n) \longrightarrow W^{k-\frac{1}{p},p}(\partial\Omega, Cl_n) \tag{3.3}$$

is the trace operator with

$$\begin{aligned} &\sum_{\|\alpha\| \leq [\lambda+\frac{1}{p}]} \int_\Omega |D^\alpha \tau f|^p dx + \sum_{\|\alpha\| = [\lambda+\frac{1}{p}]} \int_\Omega \int_\Omega \frac{|D^\alpha \tau f(x) - D^\alpha \tau f(y)|^p}{|x-y|^{n+\{\lambda+\frac{1}{p}\}p}} dxdy \\ \leq\ & \delta^p \left(\sum_{\|\alpha\| \leq [\lambda]} \int_{\partial\Omega} |D^\alpha f|^p dx + \sum_{\|\alpha\| = [\lambda]} \int_{\partial\Omega} \int_{\partial\Omega} \frac{|D^\alpha f(x) - D^\alpha f(y)|^p}{|x-y|^{n-1+\{\lambda+\frac{1}{p}\}p}} d\sigma_x d\sigma_y \right) \end{aligned}$$

Proposition 4. *The composition $\xi_{\partial\Omega} \circ \tau$ preserves regularity of a function in a Sobolev space.*

Proof. Indeed, τ makes a function to loose a regularity fractional exponent of $\frac{1}{p}$ when taken along the boundary of the domain. But the boundary or *Feuter* integral $\xi_{\partial\Omega}$ augments the regularity exponent of a function defined on the boundary by an exponent of $\frac{1}{p}$.

Therefore the composition operator $\xi_{\partial\Omega} \circ \tau$ preserves or fixes the regularity exponent of a function in a Sobolev space.

□

Proposition 5. *(Borel-Pompeiu)*
Let $f \in W^{k,p}(\Omega, Cl_n)$. Then

$$f = \xi_{\partial\Omega} \tau f + \zeta_\Omega D f$$

Corollary 1. *(i) If $f \in W_0^{k,p}(\Omega, Cl_n)$, then*

$$f = \zeta_\Omega D f$$

That is D is a right inverse for ζ_Ω and ζ_Ω is a left inverse for D over traceless spaces.

(ii) If f is monogenic function over Ω, then

$$f = \xi_{\partial\Omega}\tau f$$

Therefore monogenic functions are always Cauchy transforms of their traces over the boundary.

4. Elliptic First and Second Order BVPs

Proposition 6. *Let $f \in W^{k-1,p}(\Omega, Cl_n)$ for $k \geq 1$. Then the first order elliptic BVP:*

$$\begin{cases} Du = f \ \ in\ \Omega \\ \tau u = g \ on\ \partial\Omega \end{cases} \tag{4.1}$$

has a solution $u \in W^{k,p}(\Omega, Cl_n)$ given by

$$u(x) = \xi_{\partial\Omega} g + \zeta_\Omega f$$

Proof. The proof follows from the Borel-Pompeiu relation. As to where exactly u and g belong, we make the argument : f is in $W^{k-1,p}(\Omega, Cl_n)$ and hence from the mapping property of D, we have u to be a function in $W^{k,p}(\Omega, Cl_n)$.

Also from the mapping property of the trace operator τ we have

$$\tau u = u|_{\partial\Omega} = g \in W^{k-\frac{1}{p},p}(\partial\Omega, Cl_n)$$

□

Proposition 7. *The solution $u \in W^{k,p}(\Omega, Cl_n)$ has a norm estimate :*

$$\begin{aligned} \|u\|_{W^{k,p}(\Omega,Cl_n)} \quad \leq \quad & \gamma_1 \left(\sum_{\|\alpha\|\leq k-1} \int_{\partial\Omega} |D^\alpha g|^p d\sigma x + \sum_{\|\alpha\|=k-1} \int_{\partial\Omega}\int_{\partial\Omega} \frac{|D^\alpha g(x) - D^\alpha g(y)|^p}{|x-y|^{n+p-2}} d\sigma_x d\sigma_y \right)^{\frac{1}{p}} \\ & + \gamma_2 \left(\sum_{\|\alpha\|=k-1} \int_{\partial\Omega} |f|^p dx \right)^{\frac{1}{p}} \end{aligned}$$

where γ_1, γ_2 are constants the depend on p,n and Ω.

Proof. First let us determine regularity exponents of

$$g \in W^{k-\frac{1}{p},p}(\partial\Omega, Cl_n)$$

For the regularity index $k - \frac{1}{p}$ the integer part is

$$[k - \frac{1}{p}] = k - 1$$

and the fractional part is

$$\{k-\frac{1}{p}\}=1-\frac{1}{p}$$

Besides $\dim(\partial\Omega)=n-1$. From the mapping properties of D, ζ_Ω, τ and $\xi_{\partial\Omega}$, we have

$$u\in W^{k,p}(\Omega,Cl_n)$$

and

$$\tau u=g\in W^{k-\frac{1}{p},p}(\partial\Omega,Cl_n)$$

Therefore the solution u given by:

$$u(x)=\xi_{\partial\Omega}g+\zeta_\Omega f$$

has norm estimate

$$\begin{aligned}
\|u\|_{W^{k,p}(\Omega,Cl_n)} &= \|\xi_{\partial\Omega}g+\zeta_\Omega f\|_{W^{k,p}(\Omega,Cl_n)}\\
&\leq \|\xi_{\partial\Omega}g\|_{W^{k,p}(\Omega,Cl_n)}+\|\zeta_\Omega f\|_{W^{k,p}(\Omega,Cl_n)}\\
&\leq \gamma_1\|g\|_{W^{k-\frac{1}{p},p}(\partial\Omega,Cl_n)}+\gamma_2\|f\|_{W^{k-1,p}(\Omega,Cl_n)}\\
&= \gamma_1\left(\sum_{\|\alpha\|\leq k-1}\int_{\partial\Omega}|D^\alpha g|^p d\sigma x+\sum_{\|\alpha\|=k-1}\int_{\partial\Omega}\int_{\partial\Omega}\frac{|D^\alpha g(x)-D^\alpha g(y)|^p}{|x-y|^{n-1+\{k-\frac{1}{p}\}p}}d\sigma_x d\sigma_y\right)^{\frac{1}{p}}\\
&\quad +\gamma_2\left(\sum_{\|\alpha\|=k-1}\int_{\partial\Omega}|f|^p dx\right)^{\frac{1}{p}}\\
&= \gamma_1\left(\sum_{\|\alpha\|\leq k-1}\int_{\partial\Omega}|D^\alpha g|^p d\sigma x+\sum_{\|\alpha\|=k-1}\int_{\partial\Omega}\int_{\partial\Omega}\frac{|D^\alpha g(x)-D^\alpha g(y)|^p}{|x-y|^{n+p-2}}d\sigma_x d\sigma_y\right)^{\frac{1}{p}}\\
&\quad +\gamma_2\left(\sum_{\|\alpha\|=k-1}\int_{\partial\Omega}|f|^p dx\right)^{\frac{1}{p}}
\end{aligned}$$

The constants γ_1 and γ_2 are from the mapping properties of $\xi_{\partial\Omega},\zeta_\Omega$ and τ. □

Proposition 8. *Let $f\in W^{k,p}(\Omega,Cl_n)$. Then the second order elliptic BVP*

$$\left\{\begin{array}{c} -\Delta u=f \quad in\ \Omega\\ \tau Du=g_1 \quad on\ \partial\Omega\\ \tau u=g_2 \quad on\ \partial\Omega\end{array}\right. \tag{4.2}$$

has a solution given by

$$u=\xi_{\partial\Omega}(g_2)+\zeta_\Omega\xi_{\partial\Omega}(g_1)+\zeta_\Omega\circ\zeta_\Omega(f)$$

in $W^{k+2,p}(\Omega)$ with

$$g_1\in W^{k+1-\frac{1}{p},p}(\partial\Omega),\quad g_2\in W^{k+2-\frac{1}{p},p}(\partial\Omega)$$

Proof. As $f \in W^{k,p}(\Omega, Cl_n)$, the solution u is in the Sobolev space $W^{k+2,p}(\Omega)$ and hence

$$\tau u = g_2 \in W^{k+2-\frac{1}{p},p}(\partial\Omega)$$

But then Du is in $W^{k+1,p}(\Omega)$ and hence

$$\tau Du = g_1$$

is in the Slobodeckij space $W^{k+1-\frac{1}{p},p}(\partial\Omega)$.

The solution u of the BVP is obtained by repeated application of the Borel-Pompeiu formula by writing the Laplacian Δ as $-D^2$.

Now let us first determine the integer and fractional parts of indices $k+2-\frac{1}{p}$ and $k+1-\frac{1}{p}$ as

$$\begin{aligned}
[k+2-\frac{1}{p}] &= k+1, \quad \{k+2-\frac{1}{p}\} = 1-\frac{1}{p} \\
[k+1-\frac{1}{p}] &= k, \quad \{k+1-\frac{1}{p}\} = 1-\frac{1}{p}
\end{aligned}$$

Therefore from the properties of the mappings studied above, we have a norm estimate of the solution u in $W^{k+2,p}(\Omega)$ in terms of norms of f, g_1 and g_2 as follow:

$$\begin{aligned}
\|u\|_{W^{k+2,p}(\Omega)} &= \|\xi_{\partial\Omega}(g_2) + \zeta_\Omega \xi_{\partial\Omega}(g_1) + \zeta_\Omega \circ \zeta_\Omega(f)\|_{W^{k+2,p}(\Omega)} \\
&\leq \gamma_1 \left(\sum_{\|\alpha\| \leq k+1} \int_{\partial\Omega} |D^\alpha g_2|^p d\sigma_x + \sum_{\|\alpha\|=k+1} \int_{\partial\Omega}\int_{\partial\Omega} \frac{|D^\alpha g_2(x) - D^\alpha g_2(y)|^p}{|x-y|^{n+p-2}} d\sigma_x d\sigma_y \right)^{\frac{1}{p}} \\
&\quad + \gamma_2 \left(\sum_{\|\alpha\| \leq k} \int_{\partial\Omega} |D^\alpha g_1|^p d\sigma_x + \sum_{\|\alpha\|=k} \int_{\partial\Omega}\int_{\partial\Omega} \frac{|D^\alpha g_1(x) - D^\alpha g_1(y)|^p}{|x-y|^{n+p-2}} d\sigma_x d\sigma_y \right)^{\frac{1}{p}} \\
&\quad + \gamma_3 \left(\sum_{\|\alpha\| \leq k} \int_{\partial\Omega} |D^\alpha f|^p dx \right)^{\frac{1}{p}}
\end{aligned}$$

for some positive constants γ_1, γ_2 and γ_3 that depend on p, n, Ω □

Proposition 9. *For the BVP* (4.1) *there exist positive constants* c, γ_1 *and* γ_2 *such that the solution* $u \in W^{k,2n}(\Omega)$ *satisfies the norm estimate:*

$$
\begin{aligned}
& c^{-1}\left(\sup_{\substack{x,y\in\Omega\\ x\neq y}}\frac{|u(x)-u(y)|}{|x-y|^{\frac{1}{2}}}+\|u\|_{C(\Omega)}\right)\\
\leq\ & \gamma_1\left(\sum_{\|\alpha\|\leq k-1}\int_{\partial\Omega}|D^{\alpha}g|^{2n}d\sigma x+\sum_{\|\alpha\|=k-1}\int_{\partial\Omega}\int_{\partial\Omega}\frac{|D^{\alpha}g(x)-D^{\alpha}g(y)|^{2n}}{|x-y|^{n+p-2}}d\sigma_x d\sigma_y\right)^{\frac{1}{2n}}\\
& +\gamma_2\left(\sum_{\|\alpha\|=k-1}\int_{\partial\Omega}|f|^{2n}dx\right)^{\frac{1}{2n}}
\end{aligned}
$$

Proof. From the Sobolev embeding theorems, if $p>n$, then

$$W^{k,p}(\Omega)\hookrightarrow C^{0,\lambda}(\Omega)$$

for $0<\lambda\leq 1-\frac{n}{p}$.

But then for $p=2n$, we have $0<\lambda\leq\frac{1}{2}$ and thuerefore the solution u which is in $W^{k,2n}(\Omega)$ is contained in Hölder spaces $C^{0,\lambda}(\Omega)$.

Thus $\exists\ c=c(p,n,\Omega)>0$ such that

$$c^{-1}\|u\|_{C^{0,\lambda}(\Omega)}\leq\|u\|_{W^{k,2n}(\Omega)}$$

That is

$$
\begin{aligned}
& c^{-1}\left(\sup_{\substack{x,y\in\Omega\\ x\neq y}}\frac{|u(x)-u(y)|}{|x-y|^{\lambda}}+\|u\|_{C(\Omega)}\right)\\
\leq\ & \|u\|_{W^{k,2n}(\Omega)}\\
\leq\ & \gamma_1\left(\sum_{\|\alpha\|\leq k-1}\int_{\partial\Omega}|D^{\alpha}g|^{2n}d\sigma x+\sum_{\|\alpha\|=k-1}\int_{\partial\Omega}\int_{\partial\Omega}\frac{|D^{\alpha}g(x)-D^{\alpha}g(y)|^{2n}}{|x-y|^{n+p-2}}d\sigma_x d\sigma_y\right)^{\frac{1}{2n}}\\
& +\gamma_2\left(\sum_{\|\alpha\|=k-1}\int_{\partial\Omega}|f|^{2n}dx\right)^{\frac{1}{2n}}
\end{aligned}
$$

Choosing $\lambda=\frac{1}{2}$, we have the required result. □

References

[1] S. Bernstein, *Operator Calculus for Elliptic Boundary Value Problems in Unbounded Domains*, Zeitschrift fur Analysis Und ihre Anwendungen Vol.10 (1991) 4, 447-460.

[2] F. Brackx, R. Delanghe and F. Sommen, *Clifford Analysis*, Research Notes in Mathematics No.76, Pitman , London 1982.

[3] Di. Nezza et al., *Hitchhiker's guide to the fractional Sobolev spaces*, Bull. sci. math. (2012)

[4] K. Gürlebeck, U. Kähler, J. Ryan and W. Spröessig, *Clifford Analysis Over Unbounded Domains*, Adv. in Appl. Mathematics 19(1997), 216-239.

[5] Dejenie A. Lakew, *$W^{2,k}-$Best Approximation of a $\gamma-$Regular Function*, Journal of Applied Analysis, Vol. 13, No. 2 (2007) pp. 259-273.

[6] Dejenie A. Lakew and John Ryan, *Clifford Analytic Complete Function Systems for Unbounded Domains*, Math. Meth. in the Appl. Sci. 2002;25;1527-1539 (with John Ryan).
[7] ___, *Complete Function Systems and Decomposition Results Arising in Clifford Analysis*, Computational Methods and Function Theory, CMFT No. 1(2002) 215-228 (with John Ryan).
[8] Evans, Lawrence, *Partial Differential Equations*, Graduate Studies in Mathematics, Vol. 19, AMS, 1998
[9] S.G. Mikhlin, S. Prossdorf, *Singular Integral Operators*, Academic Verlag, Berlin (1980).
[10] John Ryan, *Intrinsic Dirac Operators in C^n*, Advances in Mathematics 118, 99-133(1996).
[11] K.T. Smith, *Primier of Modern Analysis*, Undergraduate Texts in Mathematics, Springer Verlag, New York (1983).
[12] H. Triebel, *Interpolation Theory, Function Spaces, Differential Operators*, North-Holland Mathematical Library, 1978.

John Tyler Community College, Department of Mathematics
E-mail address: dlakew@jtcc.edu
URL: http://www.jtcc.edu